


# Univariate Time Series Analysis

Kevin Sheppard

<https://kevinsheppard.com/teaching/mfe/>



$$w = \eta(R\beta - r)' [R\hat{\Sigma}_{XX}^{-1} \hat{S} \hat{\Sigma}_{XX}^{-1} R']^{-1} (R\hat{\beta} - r) \xrightarrow{d} \chi_n^2$$

$$g(e) = \frac{1}{T} \sum_{t=1}^T K\left(\frac{\hat{e}_t - e}{h}\right)$$

$$\hat{\Sigma} = \begin{bmatrix} \Sigma_{11} & \Sigma_{12} \\ \Sigma_{21} & \Sigma_{22} \end{bmatrix}$$


$$kS = \max_{\tau} \left| \sum_{i=1}^{\tau} I_{[u_i < \frac{\tau}{2}]} - \frac{\tau}{2} \right| \quad \sqrt{n}(\hat{S} - S) \xrightarrow{d} N\left(\mathbf{0}, \mathbf{1} - \frac{\mu_3}{\sigma^3} + \frac{\mu^2(\mu_4 - \sigma^4)}{4\sigma^6}\right)$$

$$\mu_r \equiv E[(X - \mu)^r] = \int_{-\infty}^{\infty} (x - \mu)^r f(x) dx$$

$$\Delta y_t = \phi_0 + \delta_1 t + \gamma y_{t-1} + \sum_{p=2}^p \phi_p \Delta y_{t-p} + \epsilon_t \quad AIC = \ln \hat{\sigma}^2 + \frac{2k}{n}$$

$$t = \frac{\sqrt{n}(R\hat{\theta} - r)}{\sqrt{RG^{-2}\Sigma(G^{-2})'R'}} \xrightarrow{d} N(\mathbf{0}, \mathbf{1}) \quad BIC = \ln \hat{\sigma}^2 + k \frac{\ln n}{n}$$

$$\frac{\mu_4}{(\sigma^2)^2} = \frac{E[(X - E[X])^4]}{E[(X - E[X])^2]^2} = E[z^4] \quad N(\mu_1 + \beta'(x_2 - \mu_2), \Sigma_{11} - \beta' \Sigma_{22} \beta)$$




$$\hat{f}(x; \rho) = \rho^x (1 - \rho)^{1-x}, \rho \geq 0$$

$$\hat{f}(\rho|x) \propto \rho^x (1 - \rho)^{1-x} \times \frac{\rho^{\alpha-1} (1 - \rho)^{\beta-1}}{B(\alpha, \beta)}$$


$$E\left[\beta \left(\mathbf{1} + r_{i,t+1}\right) \left(\frac{U'(c_{t+1})}{U'(c_t)}\right) - \mathbf{1}\right] z_t = 0$$

$$Y_t = \delta + Y_{t-1} + \epsilon_t$$


$$\sqrt{T}(R(\hat{\theta}) - R(\theta_0)) \xrightarrow{d} N\left(\mathbf{0}, \frac{\partial R(\theta_0)}{\partial \theta'} \Sigma \frac{\partial R(\theta_0)}{\partial \theta}\right)$$



$$\text{argmin}_{\beta} (y - X\beta)'(y - X\beta) + \lambda \sum_{j=1}^k |\beta_j|$$

$$\sqrt{n} \frac{y_n - \mu_n}{\sigma_n} \xrightarrow{d} N(\mathbf{0}, \mathbf{1})$$



$$Power(\mu_0, \mu_1, \sigma, \alpha, N) = \Phi\left(\frac{c_0 - \mu_1}{\frac{\sigma}{\sqrt{n}}}\right) + 1 - \Phi\left(\frac{c_0 - \mu_0}{\frac{\sigma}{\sqrt{n}}}\right)$$

$$-2X'(y - X\beta) = -2X'\epsilon = 0 \quad \beta \approx \frac{\partial Y_i}{\partial X_i} X_i = E_{y_i, x}$$


$$A \cap B = \{x : x \in A \text{ and } x \in B\}$$

$$\frac{\partial l(\theta; y)}{\partial \mu} = \sum_{i=1}^n \frac{(y_i - \mu)}{\sigma^2}$$

$$c(u_1, u_2, \dots, u_k) = \frac{\partial^k C(u_1, u_2, \dots, u_k)}{\partial u_1 \partial u_2 \dots \partial u_k}$$




$$\frac{\partial l(\theta; y)}{\partial \sigma^2} = -\frac{n}{2\sigma^2} + \frac{1}{2} \sum_{i=1}^n \frac{(y_i - \mu)^2}{\sigma^4}$$

$$\hat{f}(x_1|x_2) = \frac{\int_B \hat{f}(x_1, x_2) dx_2}{\int_B \hat{f}_2(x_2) dx_2}$$

$$z_t = \Upsilon z_{t-1} + \xi_t$$

$$\sigma_t^2 = \omega + \alpha V_{t-1}^2 + \beta \sigma_{t-1}^2$$

$$\Sigma_t = CC' + AA' \odot \epsilon_{t-1} \epsilon_{t-1}' + BB' \odot \Sigma_{t-1}$$



$$\text{VaR}_{t+1} = -\mu - \sigma_{t+1} G_{CF}^{-1}(\alpha)$$

$$\mathcal{J} = E\left[\frac{\partial l(y; \psi)}{\partial \psi} \frac{\partial l(y; \psi)}{\partial \psi'}\right]$$

## Modules

### Overview

- Key Concepts in Time Series Analysis
- Model Components
- Deterministic Processes: Trends and Seasonality
- Cyclical Processes: Autoregressive Moving-Average Processes
- Properties of ARMA Processes
- Autocorrelations and Partial Autocorrelations
- Parameter Estimation
- Model Building and Diagnostics
- Forecasting and Forecast Evaluation
- Cyclical Seasonality and Seasonal Differencing
- Random Walks and Unit Roots
- Non-linear Models for the mean

## Course Structure

- Course presented through two overlapping channel:
  1. In-person lectures
  2. Notes that accompany the lecture content
    - ▷ Read before or after the lecture or when necessary for additional background
- Slides are primary – material presented during lectures is examinable
- Notes are secondary and provide more background for the slides
- Slides are derived from notes so there is a strong correspondence

3

## Monitoring Your Progress

- Self assessment
  - ▶ Review questions in printer-friendly version of slides
    - ▷ Self-assessment
  - ▶ Multiple choice questions on Canvas made available each week
    - ▷ Answers available immediately
  - ▶ Long-form problem distributed each week
    - ▷ Answers presented in a subsequent class
- Marked Assessment
  - ▶ Empirical projects applying the material in the lectures
  - ▶ Each empirical assignment will have a written and code component

4

# Stochastic Processes

## Definition (Stochastic Process)

A stochastic process is a collection of random variables  $\{Y_t\}$  defined on a common probability space indexed by a set  $\mathcal{T}$  usually defined as  $\mathbb{N}$  for discrete time processes or  $[0, \infty)$  for continuous time processes.

**Basic Example:** An i.i.d. time series

$$Y_t \stackrel{\text{i.i.d.}}{\sim} N(0, 1)$$

5

## More Complex Examples

### ■ Random Walk

$$Y_t = Y_{t-1} + \epsilon_t, \quad \epsilon_t \stackrel{\text{i.i.d.}}{\sim} N(0, \sigma^2)$$

### ■ ARMA(1,1)

$$Y_t = \phi_1 Y_{t-1} + \theta \epsilon_{t-1} + \epsilon_t$$

► Series focuses on ARMA

### ■ GARCH(1,1)

$$Y_t \sim N(0, \sigma_t^2)$$

$$\sigma_t^2 = \omega + \alpha Y_{t-1}^2 + \beta \sigma_{t-1}^2$$


► GARCH and other non-linear processes later

### ■ Ornstein-Uhlenbeck Process

$$Y(t) = e^{-\beta t} Y(0) + \sigma \int_0^t e^{-\beta(t-s)} dW(s)$$


6

## The Default Premium




7

## Curvature of Yield Curve




8

# Industrial Production



# Housing Starts



# Review

## Stochastic Processes

### Key Concepts

#### Stochastic Process

### Questions

- What are the requirements for a sequence of random variables to be a stochastic process?
- Are cross-sectional random variables indexed by  $i$  a stochastic process?
- Are the observations of stochastic processes always regularly spaced in time?

## Autocovariance

### Definition (Autocovariance)

The autocovariance of a covariance stationary scalar process  $\{Y_t\}$  is defined

$$\gamma_s = \mathbb{E}[(Y_t - \mu)(Y_{t-s} - \mu)]$$

where  $\mu = \mathbb{E}[Y_t]$ . Note that  $\gamma_0 = \mathbb{E}[(Y_t - \mu)(Y_t - \mu)] = \mathbb{V}[Y_t]$ .

- Covariance of a process at different points in time
- Otherwise identical to usual covariance

# Stationarity

The future resembles the past

## Key concept

- Stationarity is a statistically meaningful form of regularity
- First type:

## Definition (Covariance Stationarity)

A stochastic process  $\{Y_t\}$  is covariance stationary if

$$\begin{aligned} E[Y_t] &= \mu && \text{for } t = 1, 2, \dots \\ V[Y_t] &= \sigma^2 < \infty && \text{for } t = 1, 2, \dots \\ E[(Y_t - \mu)(Y_{t-s} - \mu)] &= \gamma_s && \text{for } t = 1, 2, \dots, s = 1, 2, \dots, t - 1 \end{aligned}$$

- *Unconditional* mean, variance and autocovariance do *not* depend on time

12

# Stationarity

Second type (stronger):

## Definition (Strict Stationarity)

A stochastic process  $\{Y_t\}$  is strictly stationary if the joint distribution of  $\{Y_t, Y_{t+1}, \dots, Y_{t+h}\}$  only depends only on  $h$  and not on  $t$ .

- *Entire joint distribution* does not depend on time.
- Examples of stationary time series:
  - ▶ i.i.d. : Always strict, covariance if  $\sigma^2 < \infty$
  - ▶ i.i.d. sequence of  $t_2$  random variables, strict only
  - ▶ Multivariate normal, both
  - ▶ AR(1):  $Y_t = \phi_1 Y_{t-1} + \epsilon_t$ , covariance if  $|\phi_1| < 1$  and  $V[\epsilon_t] < \infty$ , strict is  $\epsilon_t$  is i.i.d.
  - ▶ ARCH(1):  $Y_t \sim N(0, \sigma_t^2), \sigma_t^2 = \omega + \alpha Y_{t-1}^2$  both if  $\alpha < 1$ .

13

# Nonstationarity defined

- Any series which is not stationary is nonstationary
- Four major types
  - ▶ Seasonality
    - ▷ Only slightly problematic
    - ▷ Can often be analyzed using standard tools and Box-Jenkins
  - ▶ Deterministic trends: growth over time
    - ▷ Linear
    - ▷ Polynomial
    - ▷ Exponential
  - ▶ Random walks or unit roots
  - ▶ Structural breaks

14

## What processes are not stationary?

### Nonstationary time series

- Seasonalities, Diurnality, Hebdomadality:  $Y_t = \mu + \beta I_{[\text{Quarter}(t) = Q1]} + \epsilon_t$ 
  - ▶  $E[Y_t]$  is different in Q1 than in other quarters
- Time trends:  $Y_t = t + \epsilon_t$ 
  - ▶  $E[Y_t] = t$
- Random walks:  $Y_t = Y_{t-1} + \epsilon_t$ 
  - ▶  $V[Y_t] = t\sigma^2$
- Processes with structural breaks:  $Y_t = \mu_1 + \epsilon_t$  if  $t < 1974$ ,  $Y_t = \mu_2 + \epsilon_t$ ,  $t \geq 1974$ .
  - ▶  $E[Y_t] = \mu_1 + (\mu_2 - \mu_1)(1 - I_{t < 1974})$

15

# Review

## Stationarity and Ergodicity

### Key Concepts

Covariance Stationarity, Strict Stationarity

### Questions

- Why is stationarity important when modeling and forecasting a time series?
- What is the difference between strict and covariance stationarity?
- What are the four main sources of non-stationarity in a time series?

### Problems

1. Why are the two processes below non-stationary when  $\epsilon_t \stackrel{\text{i.i.d.}}{\sim} N(0, \sigma^2)$ ?
  - a.  $Y_t = 0.3t + \epsilon_t$
  - b.  $Y_t = 0.7 + 0.2I_{[t > 2020]} + \epsilon_t$ .

# White noise

## Essential Building Block of Time Series

### Definition (White Noise)


A process  $\{\epsilon_t\}$  is known as white noise if

$$\begin{aligned} \mathbb{E}[\epsilon_t] &= 0 && \text{for } t = 1, 2, \dots \\ \mathbb{V}[\epsilon_t] &= \sigma^2 < \infty && \text{for } t = 1, 2, \dots \\ \mathbb{E}[\epsilon_t \epsilon_{t-j}] &= 0 && \text{for } t = 1, 2, \dots, j \neq 0 \end{aligned}$$


- Not necessarily independent
  - ▶ ARCH(1) process  $Y_t \sim N(0, \sigma_t^2)$ ,  $\sigma_t^2 = \omega + \alpha Y_{t-1}^2$
  - ▶ **Variance** is dependent, mean is not

# White noise


## Gaussian




## Student's $t_3$



## GARCH



## Bernoulli



17

# Linear Time-series Processes

Standard tool of time-series analysis

- Linear time series process can always be expressed as

$$Y_t = \delta_t + Y_0 + \sum_{i=0}^t \theta_i \epsilon_{t-i}$$

- Linear in the errors
- $\delta_t$  is a purely deterministic process
- $\{\epsilon_t\}$  is a White Noise process
- Example of non-linear processes
  - GARCH(1,1)

$$Y_t \sim N(0, \sigma_t^2)$$
$$\sigma_t^2 = \omega + \alpha Y_{t-1}^2 + \beta \sigma_{t-1}^2$$

- Threshold Autoregression

$$Y_t = \phi_s Y_{t-1} + \epsilon_t, \phi_s = 1 \text{ if } L < Y_{t-1} < U \text{ otherwise } 0.9$$

18

# Component View of a Time Series

$$Y_t = \underbrace{\text{Trend} + \text{Seasonal} + \text{Cyclical}}_{\text{Predictable}} + \underbrace{\text{Noise}}_{\text{Unpredictable}}$$

## Trend

- Linear, Quadratic
- Exponential
  - ▶ Linear in the log
- *Deterministic* - depends on the time period  $t$

## Seasonal

- Seasonal Summies
- Fourier Series
- *Deterministic* - depends on the time period  $t$

## Cyclical

- Autoregressive Moving Average Processes
- *Stochastic* - depends on past shocks

# Deterministic trends

- Two key types
  - ▶ Polynomial

$$Y_t = \phi_0 + \delta_1 t + \delta_2 t^2 + \dots + \delta_o t^o + \epsilon_t$$

- ▶ Linear (important special case)

$$Y_t = \phi_0 + \delta_1 t + \epsilon_t$$

- ▶ Exponential

$$\ln Y_t = \phi_0 + \delta_1 t + \epsilon_t$$

- *Mean depends on time*

$$Y_t = \phi_0 + \delta_1 t + \epsilon_t \Rightarrow E[Y_t] = \phi_0 + \delta_1 t$$

# Deterministic Seasonality

## Seasonal dummy variables

$$Y_t = \sum_{j=0}^{s-1} \beta_j I_{[t \bmod s=j]} + \epsilon_t$$

## Seasonal Fourier series

$$Y_t = \sum_{j=0}^k \lambda_j \sin\left(2\pi j \frac{t}{s}\right) + \kappa_j \cos\left(2\pi j \frac{t}{s}\right) + \epsilon_t$$

- Capture seasonal patterns using fewer terms
  - ▶  $k = 2$  in monthly data
  - ▶ 4 rather than 12 parameters
- Multiple fourier terms with different  $s$  capture additional deterministic patterns
  - ▶ Electricity: day of year, day of week, hour of day

21


# Detrending

$$Y_t = \phi_0 + \delta_1 t + \dots + \delta_o t^o + \sum_{i=0}^{s-1} \beta_i I_{[t \bmod s=i]} + \sum_{j=0}^k \lambda_j \sin\left(2\pi j \frac{t}{s}\right) + \kappa_j \cos\left(2\pi j \frac{t}{s}\right) + \epsilon_t$$

- Detrended series is a stationary process
- Detrending depends only on time  $t$
- Incorporate trends with ARMA models to capture predictable component
- Parameter estimation using OLS
- **Key problem** - most trending economic time series contains *unit roots*
  - ▶ Still not stationary even after detrending
  - ▶ Alternative: transform to remove the deterministic effects
  - ▶ More later

22


## Trending Time Series



23

## Detrended Residuals

$$\hat{\epsilon}_t = Y_t - \hat{\phi}_0 - \hat{\delta}_1 t$$



24

# ARMA Processes

- Inclusive class of all linear time-series processes

## Definition (Autoregressive-Moving Average Process)

An Autoregressive Moving Average process with orders P and Q, abbreviated ARMA(P,Q), has dynamics which follow

$$Y_t = \phi_0 + \sum_{p=1}^P \phi_p Y_{t-p} + \sum_{q=1}^Q \theta_q \epsilon_{t-q} + \epsilon_t$$

where  $\epsilon_t$  is a white noise process with the additional property that  $E_{t-1}[\epsilon_t] = 0$ .

- ARMA(1,1)

$$Y_t = \phi_1 Y_{t-1} + \theta_1 \epsilon_{t-1} + \epsilon_t$$

25

## Special case: Moving Average

- ARMA family comprises two sub-classes

## Definition (Moving Average Process of Order Q)

A Moving Average process of order Q, abbreviated MA(Q), has dynamics which follow

$$Y_t = \phi_0 + \sum_{q=1}^Q \theta_q \epsilon_{t-q} + \epsilon_t$$

where  $\epsilon_t$  is white noise series with the additional property that  $E_{t-1}[\epsilon_t] = 0$ .

- 1<sup>st</sup> order Moving Average (MA(1))

$$Y_t = \phi_0 + \theta_1 \epsilon_{t-1} + \epsilon_t$$

- Simplest non-degenerate time series process

26

# Special cases of ARMA processes: Autoregression

- Other sub-class of ARMA

## Definition (Autoregressive Process of Order $P$ )

An Autoregressive process of order  $P$ , abbreviated AR( $P$ ), has dynamics which follow

$$Y_t = \phi_0 + \sum_{p=1}^P \phi_p Y_{t-p} + \epsilon_t$$

where  $\epsilon_t$  is white noise series with the additional property that  $E_{t-1}[\epsilon_t] = 0$ .

- 1<sup>st</sup> order Autoregression (AR(1))

$$Y_t = \phi_0 + \phi_1 Y_{t-1} + \epsilon_t$$

27

## Moments and Autocovariances

$$Y_t = \phi_0 + \phi_1 Y_{t-1} + \epsilon_t$$

- Unconditional** Mean

$$E[Y_t]$$

- Unconditional** Variance

$$\gamma_0 = V[Y_t]$$

- Autocovariance

$$\gamma_s = E[(Y_t - E[Y_t])(Y_{t-s} - E[Y_{t-s}])]$$

- Conditional** Mean

$$E_t[Y_{t+1}] = E[Y_{t+1}|\mathcal{F}_t]$$

- Conditional** Variance

$$V_t[Y_{t+1}] = E_t[(Y_{t+1} - E_t[Y_{t+1}])^2]$$

28

# Review

## Linear Time Series Processes

### Key Concepts

White Noise, Linear Stochastic Process, Autoregression, Moving Average, ARMA, Conditional Moment

### Questions

- Is White Noise covariance stationary?
- Is White Noise homoskedastic?
- Is an i.i.d. sequence White Noise?
- Is an i.i.d. normal sequence White Noise?
- In what sense is a linear process *linear*?
- Why are linear processes important in the context of covariance stationary time series?
- What is the difference between a conditional and an unconditional moment?
- What is the difference between an AR and an MA model?

## How to work with ARMA processes: AR(1)

### The MA( $\infty$ ) Representation

$$Y_t = \phi_0 + \phi_1 Y_{t-1} + \epsilon_t$$

- Use backward substitution (assume  $|\phi_1| < 1$ )

$$\begin{aligned} Y_t &= \phi_0 + \phi_1 Y_{t-1} + \epsilon_t \\ &= \phi_0 + \phi_1(\phi_0 + \phi_1 Y_{t-2} + \epsilon_{t-1}) + \epsilon_t \\ &= \phi_0 + \phi_1 \phi_0 + \phi_1^2 Y_{t-2} + \phi_1 \epsilon_{t-1} + \epsilon_t \\ &= \phi_0 + \phi_1 \phi_0 + \phi_1^2(\phi_0 + \phi_1 Y_{t-3} + \epsilon_{t-2}) + \phi_1 \epsilon_{t-1} + \epsilon_t \\ &= \phi_0 \sum_{j=0}^{\infty} \phi_1^j + \sum_{i=0}^{\infty} \phi_1^i \epsilon_{t-i} \\ &= \frac{\phi_0}{1 - \phi_1} + \sum_{i=0}^{\infty} \phi_1^i \epsilon_{t-i} \end{aligned}$$

- $\lim_{s \rightarrow \infty} \sum_{i=0}^s \phi_1^i = 1/(1 - \phi_1)$

# Properties of an AR(1)

$$\begin{aligned}
 E[Y_t] &= E \left[ \frac{\phi_0}{1 - \phi_1} + \sum_{i=0}^{\infty} \phi_1^i \epsilon_{t-i} \right] \\
 &= \frac{\phi_0}{1 - \phi_1} + \sum_{i=0}^{\infty} \phi_1^i E[\epsilon_{t-i}] \\
 &= \frac{\phi_0}{1 - \phi_1} + \sum_{i=0}^{\infty} \phi_1^i 0 \\
 &= \frac{\phi_0}{1 - \phi_1}
 \end{aligned}$$

- In general AR(P):  $E[Y_t] = \frac{\phi_0}{1 - \phi_1 - \phi_2 - \dots - \phi_P}$
- Only sensible if  $\phi_1 + \phi_2 + \dots + \phi_P < 1$
- Variance can be shown in same manner
  - ▶ AR(1):  $V[Y_t] = \frac{\sigma^2}{1 - \phi_1^2}$
  - ▶ AR(P):  $V[Y_t] = \frac{\sigma^2}{1 - \rho_1\phi_1 - \rho_2\phi_2 - \dots - \rho_P\phi_P}$ 
    - ▷  $\rho$ s are autocorrelations

# Autocovariance of an AR(1)

$$\begin{aligned}
 E[(Y_t - E[Y_t])(Y_{t-s} - E[Y_{t-s}])] &= E \left[ \left( \sum_{i=0}^{\infty} \phi_1^i \epsilon_{t-i} \right) \left( \sum_{j=0}^{\infty} \phi_1^j \epsilon_{t-s-j} \right) \right] \\
 &= E \left[ \left( \underbrace{\sum_{i=0}^{s-1} \phi_1^i \epsilon_{t-i}}_{\text{After } t-s} + \underbrace{\sum_{k=s}^{\infty} \phi_1^k \epsilon_{t-k}}_{t-s \text{ and later}} \right) \left( \sum_{j=0}^{\infty} \phi_1^j \epsilon_{t-s-j} \right) \right] \\
 &= \phi_1^s \frac{\sigma^2}{1 - \phi_1^2}
 \end{aligned}$$

- Full details in notes
- The autocovariance *function*

$$\gamma_s = \phi_1^{|s|} \left\{ \frac{\sigma^2}{1 - \phi_1^2} \right\}$$

- Autocovariance declines geometrically with the lag length
- Requires  $\phi_1^2 < 1$  to exist
  - ▶ Same condition as the mean

## Stationarity of ARMA processes

- Primarily interested in covariance stationarity
- Stationarity depends on parameters of *AR* portion
- AR(0) or finite order MA: always stationary
- AR(1) or ARMA(1,Q):  $Y_t = \phi_1 Y_{t-1} + \text{MA} + \epsilon_t$ 
  - ▶  $|\phi_1| < 1$
- AR(P) or ARMA(P,Q)  $Y_t = \phi_1 Y_{t-1} + \phi_2 Y_{t-2} + \dots + \phi_P Y_{t-P} + \text{MA} + \epsilon_t$
- Rewrite  $Y_t - \phi_1 Y_{t-1} - \phi_2 Y_{t-2} - \dots - \phi_P Y_{t-P} = \text{MA} + \epsilon_t$
- Easy to determine using the characteristic equation and corresponding characteristic roots

32

## The characteristic equation

### Definition (Characteristic Equation)

Let  $Y_t$  follow a  $P^{\text{th}}$  order linear difference equation

$$Y_t = \phi_0 + \phi_1 Y_{t-1} + \phi_2 Y_{t-2} + \dots + \phi_P Y_{t-P} + x_t$$

which can be rewritten as

$$Y_t - \phi_1 Y_{t-1} - \phi_2 Y_{t-2} - \dots - \phi_P Y_{t-P} = \phi_0 + x_t$$
$$(1 - \phi_1 L - \phi_2 L^2 - \dots - \phi_P L^P) Y_t = \phi_0 + x_t$$

The characteristic equation of this process is

$$z^P - \phi_1 z^{P-1} - \phi_2 z^{P-2} - \dots - \phi_{P-1} z - \phi_P = 0$$

- Key is in the forming of the characteristic equation and its roots
- $L$  is known as “lag operator”

33

# Characteristic roots

## Definition (Characteristic Root)

Let

$$z^P - \phi_1 z^{P-1} - \phi_2 z^{P-2} - \dots - \phi_{P-1} z - \phi_P = 0$$

be the characteristic polynomial associated with some  $P^{\text{th}}$  order linear difference equation. The  $P$  characteristic roots,  $c_1, c_2, \dots, c_P$  are defined as the solution to this polynomial

$$(z - c_1)(z - c_2) \dots (z - c_P) = 0.$$

- The roots are  $c_1, c_2, \dots, c_P$
- AR(P) or ARMA(P,Q) is covariance stationary if  $|c_j| < 1$  for all  $j$
- If complex,  $|c_j| = |a_j + b_j i| = \sqrt{a^2 + b^2}$  (complex modulus)

34

## Characteristic roots example

- Difficult to determine by inspection

### Example 1

$$Y_t = .1Y_{t-1} + .7Y_{t-2} + .2Y_{t-3} + \epsilon_t$$

- Characteristic equation

$$z^3 - .1z^2 - .7z^1 - .2$$

- Roots: 1,  $-.5$ , and  $-.4 \Rightarrow$  nonstationary

### Example 2

$$Y_t = 1.7Y_{t-1} - .72Y_{t-2} + \epsilon_t$$

- Characteristic equation


$$z^2 - 1.7z^1 + .72$$

- Roots:  $.9$  and  $.8 \Rightarrow$  stationary

35

# Fitting a Basic ARMA

## YoY % change in Industrial Production



36

## Parameter Estimates

### AR2

$$Y_t = \phi_0 + \phi_1 Y_{t-1} + \phi_2 Y_{t-2} + \epsilon_t$$

### Parameter Estimates


	Estimate	s.e.	Z	p-value
$\phi_0$	0.1106	0.045	2.453	0.014
$\phi_1$	1.3187	0.017	79.114	0.000
$\phi_2$	-0.3643	0.018	-20.624	0.000
$\sigma^2$	1.3635	0.028	48.775	0.000

### Roots of Characteristic Polynomial

$c_1$	$c_2$
0.924852	0.39388

37

# Residuals



## Review

Properties or ARMA Models

### Key Concepts

Backward Substitution, Characteristic Equation, Characteristic Root

### Questions

- What role so the MA component play in determining stationarity?
- What is the key condition for stationarity of an ARMA model?
- What is complex modulus and why is it needed?

### Problems

1. Which of the models listed below are covariance stationary?
  - a.  $Y_t = 1.8Y_{t-1} - 0.8Y_{t-2} + \epsilon_t$
  - b.  $Y_t = 0.4 - 0.75Y_{t-1} - 0.25Y_{t-2} + \epsilon_t$
  - c.  $Y_t = 10 + \sum_{j=1}^{100} 0.01Y_{t-j} + \epsilon_t$
2. Write the ARMA(1,1)  $Y_t = \phi_1 Y_{t-1} + \theta_1 \epsilon_{t-1} + \epsilon_t$  as a function of  $\epsilon_t, \epsilon_{t-1}, \epsilon_{t-2}, \dots, \epsilon_{t-h}$  and  $Y_{t-h}$  using backward substitution.
3. Use backward substitution to write the model  $Y_t = -0.5\epsilon_{t-1} + \epsilon_t$  as an AR( $\infty$ ) using the relationship that  $Y_{t-1} = -0.5\epsilon_{t-2} + \epsilon_{t-1}$  implies  $\epsilon_{t-1} = Y_{t-1} + 0.5\epsilon_{t-2}$ .

# Autocorrelations and the ACF

- Autocorrelations are a **key element** of model building

## Definition (Autocorrelation)

The autocorrelation of a covariance stationary scalar process is defined

$$\rho_s = \frac{\gamma_s}{\gamma_0}$$

where  $\gamma_s = E[(Y_t - \mu)(Y_{t-s} - \mu)]$ .

- Measures the correlation of a process at different points in time
- AR(1):

$$\rho_s = \phi_1^s$$

- One of two possibilities
  - Decay geometrically if  $0 < \phi_1 < 1$
  - Oscillate and decay  $-1 < \phi_1 < 0$

# Partial Autocorrelations (PACF)

- Partial Autocorrelation is the other **key element** of model building
- More complicated than autocorrelations:
- Regression interpretation of  $s^{\text{th}}$  partial autocorrelation:

$$Y_t = \phi_0 + \phi_1 Y_{t-1} + \phi_2 Y_{t-2} + \dots + \phi_{s-1} Y_{t-s+1} + \varphi_s Y_{t-s} + \epsilon_t$$

- $\varphi_s$  is the  $s^{\text{th}}$  partial autocorrelation
  - Population (not sample) value of  $\varphi_s$
- AR(1):

$$\varphi_s = \begin{cases} \phi_1^{|s|} & \text{for } s = -1, 0, 1 \\ 0 & \text{otherwise} \end{cases}$$

- Partial autocorrelation function maps the parameters of a process to the  $s^{\text{th}}$  autocorrelation,  $\varphi(s)$

# Using the ACF and PACF to categorize processes

- ACF and PACF are useful when choosing models



Process	ACF	PACF
White Noise	All 0	All 0
AR(1)	$\rho_s = \phi_1^s$	0 beyond lag 2
AR(P)	Decays toward zero exponentially	Non-zero through lag P, 0 thereafter
MA(1)	$\rho_1 \neq 0, \rho_s = 0, s > 0$	Decays toward zero exponentially
MA(Q)	$\rho_s \neq 0, s \leq Q,$ $\rho_s = 0, s > Q$	Decays toward zero exponentially, possible oscillating
ARMA(P,Q)	Exponential Decay	Exponential Decay

# Autocorrelation for ARMA processes



ACF

PACF


White Noise




AR(1),  $\phi_1 = 0.9$




# Autocorrelation for ARMA processes



# Autocorrelation for ARMA processes



# Autocorrelation for ARMA processes



## Review

### Autocorrelation and Partial Autocorrelation

#### Key Concepts

Autocorrelation, Partial Autocorrelation

#### Questions

- What is the difference between the  $h$ -lag autocorrelation and the  $h$ -lag partial autocorrelation?
- When are the autocorrelation and partial autocorrelation always the same for any DGP?
- What shape would you expect in the ACF and PACF of an AR(3)?
- What shape would you expect in the ACF and PACF of an MA(12)?

#### Problems

1. What is the ACF and PACF of an AR(1)  $Y_t = \phi_1 Y_{t-1} + \epsilon_t$ ?
2. What is the ACF of an MA(2)  $Y_t = \theta_1 \epsilon_{t-1} + \theta_2 \epsilon_{t-2} + \epsilon_t$ ?

# Sample ACF and PACF

- Sample autocorrelations

$$\hat{\rho}_s = \frac{\sum_{t=s+1}^T Y_t^* Y_{t-s}^*}{\sum_{t=1}^T Y_t^{*2}} = \frac{\hat{\gamma}_s}{\hat{\gamma}_0}$$

►  $Y_t^* = Y_t - \bar{Y}$  where  $\bar{Y} = T^{-1} \sum_{t=1}^T Y_t$

- Some prefer the small-sample-size corrected version

$$\hat{\rho}_s = \frac{\sum_{t=s+1}^T Y_t^* Y_{t-s}^*}{\sqrt{\sum_{t=s+1}^T Y_t^{*2} \sum_{t=1}^{T-s} Y_t^{*2}}}$$

- Sample partial autocorrelations

► Run regression to estimate  $\hat{\varphi}_s$

$$Y_t = \phi_0 + \phi_1 Y_{t-1} + \phi_2 Y_{t-2} + \dots + \varphi_s Y_{t-s} + \epsilon_t$$

- More efficient ways to compute PACF using Yule-Walker (see notes)

# Testing autocorrelations and partial ACs

- Inference on autocorrelations:

$$V[\hat{\rho}_s] = T^{-1} \quad \text{for } s = 1$$

$$= T^{-1} \left( 1 + 2 \sum_{j=1}^{s-1} \hat{\rho}_j^2 \right) \quad \text{for } s > 1$$

- Standard  $t$ -stats

$$\frac{\hat{\rho}_s}{\sqrt{V[\hat{\rho}_s]}} \stackrel{A}{\sim} N(0, 1).$$

- Inference on partial autocorrelations:



$$V[\hat{\varphi}_s] \approx T^{-1}$$

- Standard  $t$ -stats

$$T^{\frac{1}{2}} \hat{\varphi}_s \stackrel{A}{\sim} N(0, 1)$$



# Autocorrelations

The Default Premium




# Autocorrelations

Monthly Housing Start Growth Rate



# Autocorrelations

## Value Weighted Market Return



50

## Testing multiple autocorrelations

- Testing multiple autocorrelations: Ljung-Box  $Q$ ,  $H_0 : \rho_1 = \dots = \rho_s = 0$

$$Q = T(T + 2) \sum_{k=1}^s \frac{\hat{\rho}_k^2}{T - k} \sim \chi_s^2$$

- **Note:** Not heteroskedasticity robust, use LM test for serial correlation

### Definition (LM test for serial correlation)

Under the null,  $E[Y_t^* Y_{t-j}^*] = 0$  for  $1 \leq j \leq s$ . The LM-test for serial correlation is constructed by defining the score vector  $\mathbf{s}_t = Y_t^* [Y_{t-1}^* \ Y_{t-2}^* \ \dots \ Y_{t-s}^*]'$ ,

$$LM = T \bar{\mathbf{s}}' \hat{\mathbf{S}}^{-1} \bar{\mathbf{s}} \xrightarrow{d} \chi_s^2$$

where  $\bar{\mathbf{s}} = T^{-1} \sum_{t=1}^T \mathbf{s}_t$ ,  $\hat{\mathbf{S}} = T^{-1} \sum_{t=1}^T \mathbf{s}_t \mathbf{s}_t'$  and  $Y_t^* = Y_t - \bar{Y}$  where  $\bar{Y} = T^{-1} \sum_{t=1}^T Y_t$ .

51

# Review

## Sample Autocorrelations and Partial Autocorrelations

### Key Concepts

Sample Autocorrelation, Sample Partial Autocorrelation, Ljung-Box Test, LM Test for Serial Correlation

### Questions

- What is the asymptotic distribution of estimated autocorrelations and partial autocorrelations?
- Where does the rule-of-thumb  $2/\sqrt{T}$  come from when plotting sample autocorrelations?
- What is the difference between the  $Q$ -test and an LM test for serial correlation?
- If you computed a sample autocorrelation in Excel using the `correl` function by copying and shifting a variable by  $h$  places, would you get the usual sample autocorrelation estimator?

## Conditional MLE

- Conditional MLE assuming distribution of  $Y_t | Y_{t-1}, \epsilon_{t-1}, Y_{t-2}, \epsilon_{t-2}, \dots$  is  $N(0, \sigma^2)$
- If  $\epsilon_{t-1}, \epsilon_{t-2}, \dots, \epsilon_{t-Q}$  are observable, identical to least squares

$$\operatorname{argmin}_{\phi, \theta} \sum_{t=P+1}^T (Y_t - \phi_0 - \phi_1 Y_{t-1} - \dots - \phi_P Y_{t-P} - \theta_1 \epsilon_{t-1} - \dots - \theta_Q \epsilon_{t-Q})^2$$

- Ignore distribution of  $Y_1, \dots, Y_P$  in fit
  - ▶ Finite sample effects, asymptotically irrelevant
- If  $\epsilon_{P-1}, \dots, \epsilon_{P-Q}$  are observable, can recursively compute  $\epsilon_P, \dots, \epsilon_T$  for a set of parameters  $\phi, \theta$
- Overcome missing initial shocks by assuming  $\epsilon_{P-1} = \dots = \epsilon_{P-Q} = 0$

# Ordinary Least Squares

- If  $Q = 0$ , conditional MLE simplifies

$$\operatorname{argmin}_{\phi} \sum_{t=P+1}^T (Y_t - \phi_0 - \phi_1 Y_{t-1} - \dots - \phi_P Y_{t-P})^2$$

- Conditional MLE is identical to OLS
- Inference is identical
- Use classical or White's covariance estimator as appropriate
- Can also incorporate deterministic terms such as time trends while maintaining simplicity of OLS

# Exact MLE

- Define the vector of data

$$\mathbf{y} = [Y_1, Y_2, \dots, Y_{T-1}, Y_T]'$$

- $\Gamma$  be the  $T$  by  $T$  covariance matrix of  $\mathbf{y}$

$$\Gamma = \begin{bmatrix} \gamma_0 & \gamma_1 & \gamma_2 & \gamma_3 & \dots & \gamma_{T-2} & \gamma_{T-1} \\ \gamma_1 & \gamma_0 & \gamma_1 & \gamma_2 & \dots & \gamma_{T-3} & \gamma_{T-2} \\ \gamma_2 & \gamma_1 & \gamma_0 & \gamma_1 & \dots & \gamma_{T-4} & \gamma_{T-3} \\ \vdots & \vdots & \vdots & \vdots & \dots & \vdots & \vdots \\ \gamma_{T-2} & \gamma_{T-3} & \gamma_{T-4} & \gamma_{T-5} & \dots & \gamma_0 & \gamma_1 \\ \gamma_{T-1} & \gamma_{T-2} & \gamma_{T-3} & \gamma_{T-4} & \dots & \gamma_1 & \gamma_0 \end{bmatrix}$$

- The joint likelihood of  $\mathbf{y}$

$$f(\mathbf{y} | \phi, \theta, \sigma^2) = (2\pi)^{-\frac{T}{2}} |\Gamma|^{-\frac{T}{2}} \exp\left(-\frac{\mathbf{y}'\Gamma^{-1}\mathbf{y}}{2}\right)$$

- Log-likelihood

$$l(\phi, \theta, \sigma^2; \mathbf{y}) = -\frac{T}{2} \ln(2\pi) - \frac{T}{2} \ln |\Gamma| - \frac{1}{2} \mathbf{y}'\Gamma^{-1}\mathbf{y}$$

# Review

## Parameter Estimation

### Key Concepts

Conditional Maximum Likelihood, Exact Maximum Likelihood

### Questions

- How are missing initial innovations addressed in conditional MLE?
- What is the key advantage of exact MLE over conditional MLE?
- When does conditional MLE reduce to OLS?
- How is the autocovariance matrix computed in exact MLE?

## Model building the Box-Jenkins way

- Model building is similar to cross-section regression
- Can use same techniques
  - ▶ General to Specific or Specific to General
  - ▶ Information criteria: AIC, BIC
- Box-Jenkins is dominant methodology, 2-steps
  - ▶ Identification: Use ACF and PACF to choose model
  - ▶ Estimation: Estimate model and do diagnostic checks
- Two principles
  - ▶ Parsimony
  - ▶ Invertibility

# Strategies

- General to Specific
  - ▶ Fit largest specification
  - ▶ Drop regressor with largest p-value
  - ▶ Refit
  - ▶ Stop if all p-values indicate significance using a size of  $\alpha$ 
    - ▷  $\alpha$  is the econometrician's choice
- Specific to General
  - ▶ Fit all specifications with a single variable
  - ▶ Retain variable with smallest p-value
  - ▶ Extend this model adding on additional variables one at a time
  - ▶ Stop if the p-values of all excluded variables are larger than  $\alpha$

56

# Information Criteria

- Information Criteria
  - ▶ Akaike Information Criterion (AIC)

$$AIC = \ln \hat{\sigma}^2 + k \frac{2}{T}$$

- ▶ Schwartz (Bayesian) Information Criterion (SIC/BIC)

$$BIC = \ln \hat{\sigma}^2 + k \frac{\ln T}{T}$$

- Both have versions suitable for likelihood based estimation
- Reward for better fit: Reduce  $\ln \hat{\sigma}^2$
- Penalty for more parameters:  $k \frac{2}{T}$  or  $k \frac{\ln T}{T}$
- Choose model with smallest IC
  - ▶ AIC has fixed penalty  $\Rightarrow$  inclusion of extraneous variables
  - ▶ BIC has larger penalty if  $\ln T > 2$  ( $T > 7$ )

57

# Model Building: Specific-to-General

The Default Premium

## AR(1)

	Estimate	s.e.	Z	p-value
$\phi_0$	3.4827	1.205	2.891	0.004
$\phi_1$	0.9652	0.007	139.901	0.000

## MA(1)

	Estimate	s.e.	Z	p-value
$\phi_0$	101.2112	2.446	41.378	0.000
$\theta_1$	0.9218	0.008	118.011	0.000

# Model Building: Specific-to-General

The Default Premium

## AR(2)

	Estimate	s.e.	Z	p-value
$\phi_0$	4.5373	1.171	3.874	0.000
$\phi_1$	1.2718	0.021	61.901	0.000
$\phi_2$	-0.3169	0.020	-15.506	0.000

## ARMA(1,1)

	Estimate	s.e.	Z	p-value
$\phi_0$	5.7953	1.587	3.652	0.000
$\phi_1$	0.9423	0.009	99.314	0.000
$\theta_1$	0.3911	0.021	18.501	0.000

# Model Building: Specific-to-General

The Default Premium

## ARMA(2,1)



	Estimate	s.e.	Z	p-value
$\phi_0$	5.8678	1.631	3.597	0.000
$\phi_1$	0.8930	0.057	15.715	0.000
$\phi_2$	0.0486	0.056	0.873	0.383
$\theta_1$	0.4337	0.052	8.412	0.000

## ARMA(1,2)

	Estimate	s.e.	Z	p-value
$\phi_0$	5.5511	1.590	3.491	0.000
$\phi_1$	0.9447	0.010	96.942	0.000
$\theta_1$	0.3814	0.024	16.024	0.000
$\theta_2$	-0.0217	0.023	-0.949	0.343

# Model Building: Information Criteria

The Default Premium





# Model Diagnostics

- Important to assess whether your model “fits”
  - ▶ Are the residuals white noise?
    - ▷ Eye-ball test
    - ▷ Ljung-Box  $Q$  stat or LM serial correlation test of  $H_0 : \rho_1 = \dots = \rho_s = 0$ .
    - ▷ SACF/SPACF of the residuals
  - ▶ Are there any large outliers?
    - ▷ Eye-ball test
- What to do if there are problems?
  - ▶ Use SPACF/SACF to repeat Box-Jenkins and augment your model with correct dynamics to pick up problem
  - ▶ Repeat diagnostics
- Concern: Repeated testing may render critical values misleading



## Ljung-Box on Residuals

ARMA(1,1)



# LM test for Serial Correlation on Residuals

ARMA(1,1)



## Review

Model Selection

### Key Concepts

Invertibility, Parsimony, AIC, BIC

### Questions

- How are the ACF and PACF used to identify candidate models?
- How does GtS differ in an ARMA from application to a linear regression?
- Which chooses a larger model, AIC or BIC, and why?
- What property should residuals have from a well specified model?
- What use is the parsimony principle?
- What does invertibility ensure?

# The information set and the law of iterated expectations

- Information set:  $\mathcal{F}_t$
- Contains a lot of information!
  - ▶ Every time  $t$  *measurable event*
  - ▶ Observed variables: prices, returns, GDP, interest rates, FX rates
  - ▶ Functions of these
  - ▶ Excludes variables which are latent: volatility

- Conditional expectation:

$$E[Y_{t+1} | \mathcal{F}_t]$$

Conditional Variance

$$V[Y_{t+1} | \mathcal{F}_t]$$

- ▶ Shorthand  $E_t[Y_{t+1}]$  and  $V_t[Y_{t+1}]$
- Law of Iterated Expectation (LIE):

$$E_t[E_{t+1}[Y_{t+2}]] = E_t[Y_{t+2}]$$

- ▶ Monday's belief about what Tuesday's belief about Wednesday is the same as Monday's belief of Wednesday

65

## Forecasting

- A  $h$ -step ahead forecast,  $\hat{Y}_{t+h|t}$ , is designed to minimize a loss function
  - ▶ MSE:  $(Y_{t+h} - \hat{Y}_{t+h|t})^2$
  - ▶ MAD:  $|Y_{t+h} - \hat{Y}_{t+h|t}|$
  - ▶ Quad-Quad:  $\alpha_1(Y_{t+h} - \hat{Y}_{t+h|t})^2 + \alpha_2 I_{[Y_{t+h} - \hat{Y}_{t+h|t} < 0]}(Y_{t+h} - \hat{Y}_{t+h|t})^2$ 
    - ▷ Asymmetric if  $\alpha_1 \neq \alpha_2$

66

## The MSE Optimal Forecast is the conditional mean

- Let  $Y_{t+h}^* = E_t[Y_{t+h}]$
- Let  $\tilde{Y}_{t+h}$  be any other value

$$\begin{aligned} E_t[(Y_{t+h} - \tilde{Y}_{t+h})^2] &= E_t\left[\left((Y_{t+h} - Y_{t+h}^*) + (Y_{t+h}^* - \tilde{Y}_{t+h})\right)^2\right] \\ &= E_t[(Y_{t+h} - Y_{t+h}^*)^2 + 2(Y_{t+h} - Y_{t+h}^*)(Y_{t+h}^* - \tilde{Y}_{t+h}) + (Y_{t+h}^* - \tilde{Y}_{t+h})^2] \\ &= V_t[Y_{t+h}] + 2E_t[(Y_{t+h} - Y_{t+h}^*)(Y_{t+h}^* - \tilde{Y}_{t+h})] + E_t[(Y_{t+h}^* - \tilde{Y}_{t+h})^2] \\ &= V_t[Y_{t+h}] + 2(Y_{t+h}^* - \tilde{Y}_{t+h})E_t[(Y_{t+h} - Y_{t+h}^*)] + E_t[(Y_{t+h}^* - \tilde{Y}_{t+h})^2] \\ &= V_t[Y_{t+h}] + 2(Y_{t+h}^* - \tilde{Y}_{t+h}) \cdot 0 + E_t[(Y_{t+h}^* - \tilde{Y}_{t+h})^2] \\ &= V_t[Y_{t+h}] + (Y_{t+h}^* - \tilde{Y}_{t+h})^2 \end{aligned}$$

67

## Forecasting

- MSE optimal forecast for an AR(1):

$$Y_t = \phi_1 Y_{t-1} + \epsilon_t$$

$$\begin{aligned} E_t[Y_{t+1}] &= E_t[\phi_1 Y_t + \epsilon_{t+1}] \\ &= \phi_1 E_t[Y_t] + E_t[\epsilon_{t+1}] \\ &= \phi_1 Y_t + 0 \end{aligned}$$


$$\begin{aligned} E_t[Y_{t+2}] &= E_t[\phi_1 Y_{t+1} + \epsilon_{t+2}] \\ &= \phi_1 E_t[Y_{t+1}] + E_t[\epsilon_{t+2}] \\ &= \phi_1 (\phi_1 Y_t) + 0 \\ &= \phi_1^2 Y_t + 0 \end{aligned}$$

**Note:** Long-run forecast is always  $E[Y_t]$  for a covariance stationary process

68

# Forecasting

## AR(1) for M2 Growth



69

## Forecast Errors

$$\begin{aligned}V_t[Y_{t+1}] &= E_t \left[ (Y_{t+1} - E_t [Y_{t+1}])^2 \right] \\&= E_t \left[ (\phi Y_t + \epsilon_{t+1} - \phi Y_t)^2 \right] \\&= E_t [\epsilon_{t+1}^2] = \sigma^2 \text{ if homoskedastic}\end{aligned}$$


$$\begin{aligned}V_t[Y_{t+2}] &= E_t \left[ (Y_{t+2} - E_t [Y_{t+2}])^2 \right] \\&= E_t \left[ (\phi^2 Y_t + \phi \epsilon_{t+1} + \epsilon_{t+2} - \phi^2 Y_t)^2 \right] \\&= E_t \left[ (\phi \epsilon_{t+1} + \epsilon_{t+2})^2 \right] \\&= \phi^2 E_t [\epsilon_{t+1}^2] + E_t [\epsilon_{t+2}^2] = (1 + \phi^2) \sigma^2 \text{ if homoskedastic}\end{aligned}$$

**Note:** Long-run forecast error variance is always  $V[Y_t]$  for a covariance stationary process

70

# Forecast Error Autocorrelation

Recursive AR(1) for M2 Growth



71

## Review

Forecasting

### Key Concepts

Mean Square Error, Conditional Expectation

### Questions

- How is the MSE optimal forecast related to the conditional mean? What about the conditional median?
- What is the key principle for producing multi-step forecasts?
- What does the long-run forecast for a covariance stationary time series always converge to? What is the long-run variance of the error?

### Problems

1. What are the first three forecasts from the model  $Y_t = \phi_0 + \phi_1 Y_{t-1} + \theta_1 \epsilon_{t-1} + \epsilon_t$ ?
2. What are the first three forecast errors?
3. What is the variance of the first three forecast errors?

# Forecast evaluation

## Mincer-Zarnowitz regressions

- Objective Forecast Evaluation

$$Y_{t+h} = \alpha + \beta \hat{Y}_{t+h|t} + \eta_t$$

- $H_0 : \alpha = 0, \beta = 1, H_1 : \alpha \neq 0 \cup \beta \neq 1$ 
  - Use any test: Wald, LR, LM
- Can be generalized to include any variable available when the forecast was produced

$$Y_{t+h} = \alpha + \beta \hat{Y}_{t+h|t} + \gamma \mathbf{x}_t + \eta_t$$

- $H_0 : \alpha = 0, \beta = 1, \gamma = \mathbf{0}, H_1 : \alpha \neq 0 \cup \beta \neq 1 \cup \gamma_j \neq 0$
- $\mathbf{x}_t$  must be in the time  $t$  information set
- Important when working with macro data

# Mincer-Zarnwotz

## AR(1) for M2 Grwoth

### Standard Form

$$Y_{t+1} = \alpha + \beta \hat{Y}_{t+1|t} + \eta_t$$

	Estimate	s.e.	Z	p-value
$\alpha$	0.0004	0.000	0.936	0.350
$\beta$	0.8481	0.061	13.985	0.000


### Simplified Form

$$Y_{t+1} - \hat{Y}_{t+1|t} = \alpha + \gamma \hat{Y}_{t+1|t} + \eta_t$$

	Estimate	s.e.	Z	p-value
$\alpha$	0.0004	0.000	0.936	0.350
$\gamma$	-0.1519	0.061	-2.505	0.013

# Mincer-Zarnwotz

AR(1) for M2 Growth



74

## Relative evaluation: Diebold-Mariano

- Two forecasts,  $\hat{Y}_{t+h|t}^A$  and  $\hat{Y}_{t+h|t}^B$
- Two losses,  $l_t^A = (Y_{t+h} - \hat{Y}_{t+h|t}^A)^2$  and  $l_t^B = (Y_{t+h} - \hat{Y}_{t+h|t}^B)^2$ 
  - ▶ Losses do not need to be MSE
- If equally good or bad,  $E[l_t^A] = E[l_t^B]$  or  $E[l_t^A - l_t^B] = 0$
- Define  $\delta_t = l_t^A - l_t^B$

75

# Relative evaluation: Diebold-Mariano

- Implemented as a  $t$ -test that  $E[\delta_t] = 0$
- $H_0 : E[\delta_t] = 0, H_1^A : E[\delta_t] < 0, H_1^B : E[\delta_t] > 0$ 
  - ▶ Composite alternative
  - ▶ Sign indicates which model is favored

$$DM = \frac{\bar{\delta}}{\sqrt{\widehat{V}[\bar{\delta}]}} = \frac{T^{-1} \sum_{t=1}^T \delta_t}{\sqrt{\frac{\hat{\sigma}_{NW}^2}{T}}}$$

- One complication:  $\{\delta_t\}$  cannot be assumed to be uncorrelated, so a more complicated variance estimator is required
- Newey-West covariance estimator:

$$\hat{\sigma}_{NW}^2 = \hat{\gamma}_0 + 2 \sum_{l=1}^L \left[ 1 - \frac{l}{L+1} \right] \hat{\gamma}_l$$

# Implementing a Diebold-Mariano Test

$$DM = \frac{\bar{\delta}}{\sqrt{\widehat{V}[\bar{\delta}]}}$$

## Algorithm (Diebold-Mariano Test)

1. Using the two forecasts,  $\hat{Y}_{t+h|t}^A$  and  $\hat{Y}_{t+h|t}^B$ , compute  $\delta_t = l_t^A - l_t^B$
2. Run the regression
 
$$\delta_t = \beta + \eta_t$$
3. Use a Newey-West covariance estimator (`cov_type="HAC"`)
4.  $T$ -test  $H_0 : \beta = 0$  against  $H_1^A : \beta < 0$ , and  $H_1^B : \beta > 0$
5. Reject if  $|t| > C_\alpha$  where  $C_\alpha$  is the critical value for a 2-sided test using a normal distribution with a size of  $\alpha$ . If significant, reject in favor of model A if test statistic is negative or in favor of model B if test statistic is positive.

# Diebold-Mariano Testing

M2 Growth: AR(1) vs a Random Walk

## Mean Square Error

$$L(Y_{t+1}, \hat{Y}_{t+1|t}) = (Y_{t+1} - \hat{Y}_{t+1|t})^2$$

	Estimate	s.e.	Z	p-value
$\delta$	$-4.365 \times 10^{-6}$	$2.16 \times 10^{-6}$	-2.017	0.044

## Mean Absolute Error



$$L(Y_{t+1}, \hat{Y}_{t+1|t}) = |Y_{t+1} - \hat{Y}_{t+1|t}|$$

	Estimate	s.e.	Z	p-value
$\delta$	-0.0003	0.000	-2.358	0.018

- OLS on a constant using Newey-West with  $[T^{1/3}]$

# Autocorrelation of MAE $\delta_t$

M2 Growth: AR(1) vs a Random Walk



# Review

## Forecast Evaluation

### Key Concepts

Objective Forecast Evaluation, Relative Forecast Evaluation, Mincer-Zarnowitz Test, Diebold-Mariano Test, Newey-West Variance Estimator

### Questions

- What is the difference between objective and relative forecast evaluation?
- Why is a Newey-West covariance estimator used in Diebold-Mariano test?
- How is rejection of the null in a Newey-West test different from most tests?
- Why is a multi-step forecast be sensitive to a future realization of the time series between the current period and the forecast horizon?
- How is a MZ regression transformed to an Augmented MZ regression?

## The Lag Operator

- The Lag Operator is a useful tool in time series
- Simplifies expressing complex models with seasonal dynamics
- Key properties
  1.  $LY_t = Y_{t-1}$
  2.  $L^2Y_t = LY_{t-1} = L(LY_t) = Y_{t-2}$
  3.  $L^a L^b = L^{(a+b)}$
  4.  $Lc = c$  where  $c$  is a constant

## Seasonality

- Seasonality is technically a form of non-stationarity
  - ▶ Mean explicitly depends on the quarter, month, day or minute
- Three types:

### Definition (Seasonality)

Data are said to be seasonal if they exhibit a non-constant deterministic pattern on an annual basis.

### Definition (Hebdomadality)

Data which exhibit day-of-week deterministic effects are said to be hebdomadal.

### Definition (Diurnality)

Data which exhibit intra-daily deterministic effects are said to be diurnal.

81

## Seasonality

- Simpler to think of processes with seasonality as having two models
  - ▶ Short-run AR and MA dynamics
  - ▶ Seasonal AR and MA dynamics
- Model building is standard with these two goals in mind

82

# ARMA Modeling of Seasonality

## Four Components

- Observation AR

$$(1 - \phi_1 L) Y_t = \phi_0 + \epsilon_t$$

- Seasonal AR

$$(1 - \phi_s L^s) Y_t = \phi_0 + \epsilon_t$$

- Observation MA

$$Y_t = \phi_0 + (1 + \theta_1 L^1) \epsilon_t$$

- Seasonal MA

$$Y_t = \phi_0 + (1 + \theta_s L^s) \epsilon_t$$

- Combined Model

$$(1 - \phi_1 L) (1 - \phi_s L^s) Y_t = (1 + \theta_1 L^1) (1 + \theta_s L^s) \epsilon_t$$

$$Y_t = \phi_0 + \phi_1 Y_{t-1} + \phi_s Y_{t-s} - \phi_1 \phi_s Y_{t-s-1} + \theta_1 \epsilon_{t-1} + \theta_s \epsilon_{t-s} + \theta_1 \theta_s \epsilon_{t-s-1} + \epsilon_t$$

# ARMA Modeling of Seasonality


## Four Components

- Generalizes to higher orders of each term
- Known as SARIMA( $p, 0, q$ )  $\times$  ( $P, 0, Q, s$ )
- Imposes restrictions on parameters due to multiplication of terms
- Can estimate unrestricted equivalent


$$Y_t = \phi_0 + \phi_1 Y_{t-1} + \phi_s Y_{t-s} + \phi_{s+1} Y_{t-s-1} + \theta_1 \epsilon_{t-1} + \theta_s \epsilon_{t-s} + \theta_{s+1} \epsilon_{t-s-1} + \epsilon_t$$

- Can test  $H_0 : \phi_{s+1} = \phi_1 \phi_s \cap \theta_{s+1} = \theta_1 \theta_s$



# Housing Starts



# YoY Growth in Housing Starts





# YoY Growth in Housing Starts Autocorrelation



# Modeling YoY Growth in Housing Starts

AR(1) Residuals



# Modeling Housing Starts

Levels



$$\text{SARIMAX}(2, 0, 0) \times (0, 0, 1, 12)$$

	Estimate	s.e.	Z	p-value
$\phi_1$	0.6809	0.034	20.284	0.000
$\phi_2$	0.2824	0.034	8.233	0.000
$\theta_{s,12}$	-0.8795	0.017	-50.520	0.000

# Modeling Housing Starts

Levels

$$\text{SARIMAX}(2, 0, 0) \times (0, 0, 1, 12)$$



# Modeling Housing Starts

Seasonal Differencing



$$\text{SARIMAX}(1, 0, 1) \times (0, 1, 1, 12)$$

	Estimate	s.e.	Z	p-value
$\phi_1$	0.9779	0.008	127.034	0.000
$\theta_1$	-0.3129	0.033	-9.361	0.000
$\theta_{s,12}$	-0.8775	0.018	-48.079	0.000

# Modeling Housing Starts

Seasonal Differencing

$$\text{SARIMAX}(1, 0, 1) \times (0, 1, 1, 12)$$



# Modeling Housing Starts


## Seasonal Dummies

SARIMAX(2, 1, 0) with Seasonal Dummies

	Estimate	s.e.	Z	p-value
$\phi_0$	0.0002	0.004	0.046	0.964
Feb	0.0358	0.012	2.965	0.003
Mar	0.3075	0.012	24.776	0.000
Apr	0.4289	0.015	29.516	0.000
May	0.4669	0.018	26.260	0.000
Jun	0.4697	0.019	25.309	0.000
Jul	0.4328	0.019	23.265	0.000
Aug	0.4117	0.019	22.227	0.000
Sep	0.3657	0.017	21.803	0.000
Oct	0.3921	0.015	26.253	0.000
Nov	0.2169	0.013	16.943	0.000
Dec	0.0502	0.010	5.242	0.000
$\phi_1$	-0.2675	0.033	-8.114	0.000
$\phi_2$	-0.1107	0.034	-3.276	0.001

# Modeling Housing Starts

## Seasonal Dummies



# Review

## Seasonality

### Key Concepts

Seasonality, Lag Operator, SARIMA, Deterministic Trend, Exponential Trend

### Questions

- How can seasonality be modeled in an ARMA model?
- Define diurnality, hebdomadality and seasonality.
- What are seasonal determinist terms and how do they differ from seasonal AR and MA terms?
- What is an exponential trend?
- What do the orders in a SARIMA mean?
- How could a standard AR be used to model a time series with a seasonal AR component?

94

## Stochastic trends

- Stochastic trends are similar to deterministic trends
  - ▶ Dominant feature of a process

$$Y_t = \text{stochastic trend} + \text{stationary component} + \text{noise}$$

- Most common stochastic trend is a unit root
- There are others (generally non-linear)
- Removed using stochastic detrending (differencing)
  - ▶ Meaningfully different that deterministic detrending

95

## Short-run Dynamics in a Unit Root process

- Unit root processes, in the long-run, behave like random walks
- In the short run, can have stationary dynamics

$$Y_t = \phi_1 Y_{t-1} + \phi_2 Y_{t-2} + \phi_3 Y_{t-3} + \epsilon_t$$

- If this process contains a unit root,  $\phi_1 + \phi_2 + \phi_3 = 1$
- Can see the SR dynamics by differencing

$$Y_t = \phi_1 Y_{t-1} + \phi_2 Y_{t-2} + \phi_3 Y_{t-2} - \phi_3 Y_{t-2} + \phi_3 Y_{t-3} + \epsilon_t$$

$$Y_t = \phi_1 Y_{t-1} + \phi_2 Y_{t-2} + \phi_3 Y_{t-2} - \phi_3 \Delta Y_{t-2} + \epsilon_t$$

$$Y_t - Y_{t-1} = (\phi_1 + \phi_2 + \phi_3 - 1) Y_{t-1} - (\phi_2 + \phi_3) \Delta Y_{t-1} - \phi_3 \Delta Y_{t-2} + \epsilon_t$$

$$\Delta Y_t = \pi_1 \Delta Y_{t-1} + \pi_2 \Delta Y_{t-2} + \epsilon_t$$

96

## What's the problem with unit roots?

- Unit roots cause a number of problems
  - ▶ Exploding variance:  $V[Y_t] = t\sigma^2$
  - ▶ Parameter estimates converge at different rates
  - ▶ Hypothesis tests have non-standard distributions
  - ▶ No mean reversion in long-run forecasts
  - ▶ Spurious regression
- Crucial to understand whether a process is stationary or contains a unit root
- Often has large economic consequences
  - ▶ PPP
  - ▶ Covered interest rate parity
  - ▶ Carry trades

97

# Testing for unit roots

- Dickey-Fuller looks like a standard  $t$ -test

$$Y_t = \phi_1 Y_{t-1} + \epsilon_t$$

- $H_0 : \phi_1 = 1, H_1 : \phi_1 < 1$
- Impose the null

$$\begin{aligned}
 Y_t - Y_{t-1} &= \phi_1 Y_{t-1} - Y_{t-1} + \epsilon_t \\
 \Delta Y_t &= (\phi_1 - 1) Y_{t-1} + \epsilon_t \\
 \Delta Y_t &= \gamma Y_{t-1} + \epsilon_t
 \end{aligned}$$

- New  $H_0 : \gamma = 0, H_1 : \gamma < 0$
- Test with  $t$ -stat
- Augmented Dickey Fuller (ADF) captures short run dynamics as well

$$\Delta Y_t = \gamma Y_{t-1} + \rho_1 \Delta Y_{t-1} + \rho_2 \Delta Y_{t-2} + \dots + \rho_P \Delta Y_{t-P} + \epsilon_t$$

- Lags of  $\Delta Y_{t-1}$  needed to ensure  $\epsilon_t \sim WN(0, \sigma^2)$ , also reduce variance of residuals

# The problem

- $t$ -stat is no longer asymptotically normal
- Requires Dickey-Fuller distribution
  - ▶ Most software packages contain the correct critical value
- Many processes with unit roots also contain deterministic components
- Asymptotic distribution depends on choice of model:

$$\Delta Y_t = \gamma Y_{t-1} + \sum_{p=1}^P \phi_p \Delta Y_{t-p} + \epsilon_t \quad \text{(No trend)}$$

$$\Delta Y_t = \delta_0 + \gamma Y_{t-1} + \sum_{p=1}^P \phi_p \Delta Y_{t-p} + \epsilon_t \quad \text{(Constant, linear in } Y_t \text{)}$$

$$\Delta Y_t = \delta_0 + \delta_1 t + \gamma Y_{t-1} + \sum_{p=1}^P \phi_p \Delta Y_{t-p} + \epsilon_t \quad \text{(Constant, quadratic in } Y_t \text{)}$$

- More deterministic regressors lower the critical value
- Reject null of unit root if  $t$ -stat of  $\gamma$  is **negative** and below the critical value

# The Role of The Deterministic Terms

- ADF tests include deterministic terms to remove these effects from  $Y_{t-1}$
- Suppose  $Y_t$  is a pure time trend process

$$Y_t = \alpha + \beta t + \epsilon_t$$

- The differenced value is

$$\begin{aligned} \Delta Y_t &= \alpha + \beta t + \epsilon_t - \alpha - \beta(t-1) - \epsilon_{t-1} \\ &= \beta - \epsilon_{t-1} + \epsilon_t \end{aligned}$$

- ▶ MA(1) without a trend

- In an ADF with deterministic regressors

$$\Delta Y_t = \delta_0 + \delta_1 t + \gamma Y_{t-1} + \epsilon_t$$

- The deterministic terms remove deterministic components from  $Y_{t-1}$

- $\gamma$  depends on


$$\text{Cov}[\Delta Y_t, Y_{t-1} - \alpha - \beta(t-1)] = \text{Cov}[\beta - \epsilon_{t-1} + \epsilon_t, \epsilon_{t-1}] = -\sigma^2$$

- Failing to include the deterministic regressors results in  $\gamma$  that depends on

$$\text{Cov}[\Delta Y_t, Y_{t-1}] = 0$$

- ▶ Time trend dominates the other components of  $Y_{t-1}$

# The Dickey-Fuller Distributions



# Important considerations

- Unit root tests are well known for having low power
- Power = 1-Pr(type II)
  - ▶ Chance you don't reject when alternative is true
- Some suggestions
  - ▶ Use a loose model selection when choosing the number of lags of  $\Delta Y_{t-j}$ , e.g. AIC
  - ▶ Be conservative in excluding deterministic regressors.
    - ▷ Including a constant or time-trend when absent hurts power
    - ▷ Excluding a constant or time-trend when present results in **no power**
  - ▶ More powerful tests than the ADF are available: DF-GLS
  - ▶ Visually inspect the data and differenced data
  - ▶ Use a general-to-specific search
- Number of differences needed is the *order of integration*
  - ▶ Integrated of Order 1 or I(1):  $Y_t$  is nonstationary but  $\Delta Y_t$  is stationary
  - ▶ I(d):  $Y_t$  is nonstationary,  $\Delta^j Y_t$  also nonstationary when  $j < d$ ,  $\Delta^d Y_t$  is stationary

# Unit Root Testing

	ADF Statistic	p-value	Lags	Deterministic
Default	-3.866	0.002	10	c
Curvature	-4.412	0.000	19	c
ln Ind Prod	-2.186	0.211	4	c
	-1.831	0.690	6	ct
	-2.962	0.314	6	ctt
$\Delta$ ln Ind Prod	-11.945	0.000	3	c

- Lags determined using AIC
- Deterministic order increased when null is not rejected

# The Role Of Deterministics

Trend Stationary AR(1)

$$Y_t = 0.025t + 0.7Y_{t-1} + \epsilon_t$$

ADF Statistic	p-value	Lags	Deterministic
1.934	0.988	9	n
-1.146	0.696	9	c
-6.790	0.000	0	ct
-6.885	0.000	0	ctt

- Correct specification uses “ct”

104

# Seasonal Differencing

- Seasonal series should use seasonal differencing

$$\Delta_s Y_t = Y_t - Y_{t-s}$$

- Complete SARIMA( $P, D, Q$ )  $\times$  ( $P_s, D_s, Q_s, s$ ) model
  - ▶  $D$  is order of observational difference
  - ▶  $D_s$  is order of seasonal difference
  - ▶  $P$  and  $Q$  are observational AR and MA orders
  - ▶  $P_s$  and  $Q_s$  are seasonal AR and MA orders
- Special Cases
  - ▶ ARMA( $P, Q$ ):  $D = D_s = P_s = Q_s = 0$
  - ▶ ARIMA( $P, D, Q$ ):  $D_s = P_s = Q_s = 0$
  - ▶ SARMA( $P, Q$ )  $\times$  ( $P_s, Q_s, s$ ):  $D = D_s = 0$

105

# Review

## Unit Roots and Integration

### Key Concepts

Unit Root, Integrated Process,  $I(1)$ , Augmented Dickey-Fuller Test, Seasonal Difference

### Questions

- What happens if a relevant deterministic term is omitted in a ADF test?
- What is the effect of including an unnecessary deterministic in an ADF test?
- How should you decide how many lags of the differenced variable to include in an ADF test?
- When should you use seasonal differencing?
- What is the relationship between a random walk and a unit root process?
- What are the consequences of ignoring a unit root when modeling a time series?

105

## Nonlinear Models for the mean

- *Linear* time series process

$$Y_t = Y_0 + \sum_{i=0}^t \theta_i \epsilon_{t-i}$$

- Alternatives
  - ▶ Markov Switching Autoregression (MSAR)
  - ▶ Threshold Autoregression (TAR) and Self-exciting Threshold Autoregression (SETAR)
  - ▶ Many, many others
- Nonlinear models can capture different dynamics
  - ▶ *State-dependent parameters*

$$Y_t = \phi_0^{s_t} + \phi_1^{s_t} Y_{t-1} + \sigma^{s_t} \epsilon_t$$

- ▶ Models differ in how  $s_t$  evolves

106

# Markov Switching Example

- Two states,  $H$  and  $L$

$$Y_t = \begin{cases} \phi^H + \epsilon_t \\ \phi^L + \epsilon_t \end{cases}$$

- States evolve according to a 1<sup>st</sup> order Markov Chain


$$\{s_t\} = \{H, H, H, L, L, L, H, L, \dots\}$$

- Transition Probabilities


$$\begin{bmatrix} p_{HH} & p_{HL} \\ p_{LH} & p_{LL} \end{bmatrix} = \begin{bmatrix} p_{HH} & 1 - p_{LL} \\ 1 - p_{HH} & p_{LL} \end{bmatrix}$$

- $p_{HH}$  is the probability  $s_{t+1} = H$  given  $s_t = H$ .
- Model will switch between a high mean state and a low mean state
- Models like this are very flexible and nest ARMA
  - Successful in financial econometrics for asset allocation, volatility modeling, modeling series with business-cycle length patterns: GDP

# Markov Switching: i.i.d. Mixture




## Markov Switching: Symmetric Persistent




109

## Markov Switching: Asymmetric Persistent



110

# Markov Switching: Different Variances



## Review

Non-linear Time Series Models

### Key Concepts

Self-exciting Threshold Autoregression, Markov Switching Processes

### Questions

- It is always necessary to consider nonlinear models to model covariance stationary time series?
- What advantages might a nonlinear model have over a linear model when modeling a covariance stationary time series?